

Evidence Table 3. Patient/Provider Education: Asthma Self-Management Education for Adults

Abbreviations used in table:

COPD	chronic obstructive pulmonary disease	MDI/DED	metered-dose inhaler (MDI) with delivery enhancement device (DED)
DPI	dry powder inhaler	OR	odds ratio
ED	emergency department	PCP	primary care provider
FEF	forced expiratory flow	PEFR	peak expiratory flow rate
FEV₁	forced expiratory volume in 1 sec.	pMDI	pressurized metered-dose inhalers
FVC	forced vital capacity	SpO₂	saturation of oxyhemoglobin
HMO	health maintenance organization	95% CI	95% Confidence Interval
ICS	inhaled corticosteroid		
ITT	intent-to-treat analysis		

* indicates primary outcome

Evidence Table 3. Patient/Provider Education: Asthma Self-Management Education for Adults

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
A. ASTHMA SELF-MANAGEMENT EDUCATION FOR ADULTS IN CLINIC-BASED SETTINGS				
<p>Ford et al. Health outcomes among African American and Caucasian adults following a randomized trial of an asthma education program. <i>Ethn Health</i> 1997;2(4):329–339. (Agency for Health Care Policy and Research)</p>	<p>Secondary analysis of data from a randomized trial (2 EDs, 1 inner city and 1 suburban, of a large Midwestern health care system)</p>	<p>241 (241 ITT)</p>	<p><u>African American (n=163)</u> Age 18–70 yr, mean = 35.8 yr Gender 30.1% male, 69.9% female Smoking 28.2% current smokers, 58.3% exsmoker, 13.5% never smoked Other 27.6% married, 66.3% with income <\$15,000/year, 39.7% with education over 12 years, 52.8% employed/student <u>Caucasian (n=78)</u> Age 18–70 yr, mean = 40.2 yr Gender 89.2% male, 10.8% female Smoking 20.5% current smokers, 59.0% exsmoker, 20.5% never smoked Other 57.7% married, 25.6% with income <\$15,000/year, 65.6% with education over 12 years, 64.1% employed/student</p>	<p><u>African American (n=163)</u> Average ED visits due to asthma, mean = 5.9/yr Average days with limited activity due to asthma, mean = 12.2 days/yr Taking medication for asthma, 94.5% Overall health rating of good to excellent, 46.6%</p> <p><u>Caucasian (n=78)</u> Average ED visits due to asthma, mean = 5.5/yr Average days with limited activity due to asthma, mean = 10.8 days/yr Taking medication for asthma, 94.9% Overall health rating of good to excellent, 44.9%</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
<p>Klein et al. Benefit from the inclusion of self-treatment guidelines to a self-management programme for adults with asthma. <i>Eur Respir J</i> 2001;17(3): 386–394.</p> <p>(Netherlands Asthma Foundation; GlaxoWellcome; the “Stichting Astmabestrijding;” Amicon Health Care Insurance Fund)</p>	<p>Single-blind randomized, parallel group trial (outpatient clinic of a teaching hospital)</p>	<p>245 (238 at 12 months; 173 at 24 months)</p>	<p>Age 18–65 yr, mean = 44 yr</p> <p>Gender 45% male, 55% female</p> <p>Education 35% low, 38.3% medium, 26.7% high</p> <p>Smoking 53.1% nonsmokers, 35.5% exsmokers, 11.4% smokers</p>	<p>Stable, moderate-to-severe asthma</p> <p>Duration of asthma, mean = 20 yr</p> <p>Continuous use of inhaled steroids for at least 3 months, MDI equivalent 589 mcg/day</p> <p>Morning PEF, mean = 385 L/min</p> <p>FEV₁ % pred., mean = 76.5</p> <p>Positive skin prick test = 67.5%</p>
<p>Levy et al. A randomized controlled evaluation of specialist nurse education following accident and emergency department attendance for acute asthma. <i>Respir Med</i> 2000;94(9):900–908.</p> <p>(National Asthma Campaign, UK; Allen and Hanbury’s; Clement Clarke International; Astra Pharmaceuticals)</p>	<p>Randomized controlled trial (emergency department in 3 local hospitals)</p>	<p>103 (103 ITT)</p>	<p>Age >18 yr, mean = 40 yr</p> <p>Gender 38% male, 62% female</p> <p>Ethnicity Not reported</p>	<p>Attending accident and emergency departments or admitted to hospital for uncontrolled asthma</p> <p>79% nebulized in ED; 64% prescribed oral steroids; 32% admitted</p>
<p>Neri et al. Short and long-term evaluation of two structured self management programmes on asthma. <i>Monaldi Arch Chest Dis</i> 2001;56(3): 208–210.</p>	<p>Randomized controlled trial</p>	<p>40 (40)</p>	<p>Age Mean = 46.5 yr</p> <p>Gender 50% male, 50% female</p>	<p>Asthma duration at least 1 yr</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
<p>Van der Palen et al. Behavioural effect of self-treatment guidelines in a self-management program for adults with asthma. Patient Educ Couns 2001;43(2):161–169. (The Netherlands Asthma Foundation; the "Stichting Astmabestrijding Nederland"; GlaxoWellcome)</p>	<p>Randomized controlled parallel group trial</p>	<p>245 (238)</p>	<p>Age 18–65 yr, mean 44.3 yr Gender 45% male, 55% female Education 35% low, 38% medium, 27% high Smoking 53% nonsmokers, 35% exsmokers, 12% smokers</p>	<p>Stable phase of disease Inhaled steroids, mean = 594 mcg/day metered dose inhaler equivalent FEV₁ % pred., mean = 76.5 Morning PEF, mean = 384.4 l/min</p>
<p>Couturaud et al. Education and self-management: a one-year randomized trial in stable adult asthmatic patients. J Asthma 2002;39(6):493–500. (Projets Hospitaliers de Recherche Clinique 1995)</p>	<p>Multicenter, prospective, open, randomized parallel group design (outpatient clinic of 2 university hospitals)</p>	<p>72 (54)</p>	<p>Age 18–72 yr, mean = 38 yr Gender 32% male, 68% female Smoking 12% current smokers</p>	<p>Moderate to severe asthma FEV₁ % pred., mean = 84 Inhaled steroids, mean = 1562 mcg/day Long-acting beta₂-agonists = 67% 75% hospitalized in past 5 years</p>
<p>Cowie et al. Asthma in adolescents: a randomized, controlled trial of an asthma program for adolescents and young adults with severe asthma. Can Respir J 2002;9(4):253–259. (Alberta, Canada, Lung Association)</p>	<p>Randomized controlled trial (recruited from EDs in Calgary, Alberta)</p>	<p>93 (93 ITT)</p>	<p>Age 15–20 yr, mean = 17.2 yr Gender 29% male, 71% female</p>	<p>Severe asthma 26% admitted for asthma in year before study 76% using inhaled corticosteroid 56% waking with asthma, mean of 2 times/week 77% using beta₂-agonist more than once/day FEV₁ % pred., mean = 81</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
Gibson et al. Self-management education and regular practitioner review for adults with asthma (Review). Cochrane Database Syst Rev 2002;(2):CD001117. (NSW Health Cooperative Research Centre for Asthma, Australia)	Meta-analysis and Systematic Review of randomized controlled trials and controlled clinical trials	36 trials published 1986–2001 6090 participants randomized; 4593 completers; drop out rates ranged from 0% to 54%	Age Over 16 years	
Gibson et al. Limited (information only) patient education programs for adults with asthma (Review). Cochrane Database Syst Rev 2002;(2):CD001005. (Cooperative Research Centre for Asthma, Australia; Garfield Weston Foundation UK)	Meta-analysis and Systematic Review of randomized and controlled trials	12 trials published 1979–1997	Age Over 16 years	
Marabini et al. Short-term effectiveness of an asthma educational program: results of a randomized controlled trial. Respir Med 2002;96(12):993–998. (National Research Council, Italy)	Randomized controlled trial (outpatients attending asthma clinic, Italy)	77 (77)	Age ≥18 yr, mean = 51.1 yr Gender 46.8% male, 53.2% female Ethnicity Not reported Smoking 59.7% nonsmokers, 29.9% exsmokers, 10.4% current smokers Employment Status 15.6% unemployed or retired	Primary diagnosis of persistent asthma for at least 1 year: 16.9% mild, 64.9% moderate, 18.2% severe Duration of asthma, mean = 14.2 yr Stable asthma in 3 months before recruitment Atopy, 35.1%

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
<p>Perneger et al. Effect of patient education on self-management skills and health status in patients with asthma a randomized trial. Am J Med 2002; 113(1):7–14. (Swiss National Science Foundation, Berne, Switzerland)</p>	<p>Randomized controlled trial (1 university hospital)</p>	<p>131 (115)</p>	<p>Age <30 yr, 23.7% 30–44 yr, 30.5% 45–59 yr, 28.2% 60+ yr, 17.6% Gender 40% male, 60% female Smoking 34% current smokers, 26% exsmokers, 40% never smoked Other 26% university education</p>	<p>Hospitalized for asthma (76%), seen in the emergency ward, or received asthma medications while hospitalized for another reason Confirmed diagnosis of asthma Duration of asthma: <2 yr, 18%; 2–10 yr, 28%; >10 yr, 54%</p>
<p>Powell et al. Options for self-management education for adults with asthma (Review). Cochrane Database Syst Rev 2003;(1):CD004107. (Cooperative Research Centre for Asthma, Australia)</p>	<p>Meta-analysis and systematic review of randomized controlled trials</p>	<p>15 studies published 1990–2001 2460 participants randomized; dropout rates ranged from 0% to 60.3% with rate not reported for 1 study</p>	<p>Age Over age 16 years</p>	
<p>Janson et al. Effects of individual self-management education on clinical, biological, and adherence outcomes in asthma. Am J Med 2003;115(8):620–626. (National Institutes of Health)</p>	<p>Prospective, randomized controlled trial</p>	<p>68 (65)</p>	<p>Age ≥18 yr, mean = 33.5 yr Gender 45% male, 55% female Education Mean = 16 yr Smoking All nonsmoking</p>	<p>Mild-to-moderate persistent asthma All prescribed an inhaled corticosteroid of at least 400 mcg/day Duration mean = 19.5 yr FEV₁ % pred., mean = 78.5 Symptom severity score mean = 10 out of 50 Nights with symptoms, mean = 2/week Perceived asthma control score, mean = 39 out of 55 Asthma quality of life score, mean = 25.5 out of 80 Methacholine PC₂₀, mean = 0.80 mg/mL</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
<p>Thoonen et al. Self-management of asthma in general practice, asthma control and quality of life: a randomized controlled trial. <i>Thorax</i> 2003;58(1):30–36. (The Netherlands Organization for Scientific Research; ASTRAZeneca Pharmaceutica BV)</p>	<p>Nonrandomized controlled trial (practices randomized; unit of analysis was individual patient)</p>	<p>19 practices; 214 patients (193 ITT)</p>	<p>Age >16 yr, mean = 39.5 yr Gender 38% male, 62% female Smoking status 52% never smokers, 27% former smokers, 22% current smokers Employment 68% employed full-time or part-time</p>	<p>Stable asthma Duration of asthma, mean = 19.6 yr 40% with asthma attack in previous 6 months FEV₁ % pred. post-bronchodilator, mean = 91.3 Asthma Quality of Life Questionnaire score, mean = 5.5</p>
<p>Hopman et al. Outcomes of asthma education: results of a multisite evaluation. <i>Can Respir J</i> 2004;11(4): 291–297.</p>	<p>Multicenter, prospective, observational, comparative study; pre-post design (7 sites across Canada: 5 Community Asthma Care Centres, 1 community hospital, 1 academic health sciences center)</p>	<p>396 (252)</p>	<p>Age ≥16 yr; mean = 42 yr Gender 24% male, 76% female</p>	<p>All referred for education on basis of poor asthma control and/or exacerbation of asthma Action plan, 31% Inhaled corticosteroids, 90.5% Corticosteroid tablets, 18% Short-acting beta-agonist, 90.9% Long-acting beta-agonist, 11.5%</p>
<p>Magar et al. Assessment of a therapeutic education programme for asthma patients: “un soufflé nouveau.” <i>Patient Educ Couns</i> 2005;58(1):41–46. (GlaxoSmithKline)</p>	<p>Multicenter randomized controlled trial (26 pneumologists in 4 centres in France)</p>	<p>238 (193)</p>	<p>Age 18–60 yr Gender Not reported</p>	<p>Asthma diagnosed according to GINA guidelines All required ongoing inhaled corticosteroids and/or long acting beta₂-agonists In the previous month, averaged 1 daytime asthma attack per week, 1 nighttime asthma attack per week, or had to use bronchodilator once per week</p>
<p>Urek et al. Effect of educational programs on asthma control and quality of life in adult asthma patients. <i>Patient Educ and Couns</i> 2005;58(1):47–54.</p>	<p>Randomized trial (clinical site in Croatia)</p>	<p>60 (60)</p>	<p>Age mean = 43.4 yr Gender 35% male, 65% female</p>	<p>Moderate persistent asthma based on GINA guidelines criteria FVC % pred., mean = 93.9 FEV₁ % pred., mean = 68.9 FEF₅₀ % pred., mean = 55.8 PEF % pred., mean = 72.1</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
B. ASTHMA SELF-MANAGEMENT EDUCATION FOR ADULTS IN HOSPITAL-BASED SETTINGS				
George et al. A comprehensive educational program improves clinical outcome measures in inner-city patients with asthma. Arch Intern Med 1999;159(15):1710–1716. (NIH; Health Management Alternatives, Abbott Laboratories; Mary Rockefeller Foundation; Merck & Co, West Point, PA; Schering-Plough/Key Pharmaceuticals, Kenilworth, NJ)	Randomized controlled trial (patients admitted to hospital from university hospital ED)	77 (77; 50 for data from Medicaid MCO database)	Age 18–45 yr, mean = 29 yr Gender 20.8% male, 79.2% female Insurance Medicaid, 57.1%; Self-pay, 11.7%; Private, 31.2%	Admitted to the hospital from the ED with acute exacerbation
Morice & Wrench. The role of the asthma nurse in treatment compliance and self-management following hospital admission. Respir Med 2001;95(11):851–856. (Astra Pharmaceuticals)	Randomized controlled trial (large teaching hospital in United Kingdom)	80 (70 at 6 weeks; 65 at 6 months)	Age 16–72 yr, mean = 36.1 yr Gender 34% male, 66% female	Admitted to hospital with documented primary diagnosis of acute asthma
Osman et al. A randomized trial of self-management planning for adult patients admitted to hospital with acute asthma. Thorax 2002;57(10):869–874. (NHS R&D Programme in Delivery of Care in Asthma managed by the National Asthma Campaign, UK)	Randomized controlled trial (7 general medical units and the respiratory unit of a 1022-bed teaching hospital)	280 (226 at 1 month; 271 at 12 months)	Age 14–60 yr, median 29 yr Gender 34% male, 66% female Smoking 43% current smoker	Admitted to hospital with acute asthma Median admission PEF % pred., = 41.2 Inhaled steroid prescribed before admission = 79% Previous admission for acute asthma: 51% never, 45% within 10 yr, 4% within 11–20 yr Median yr since previous admission = 2

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
<p>Castro et al. Asthma intervention program prevents readmissions in high healthcare users. Am J Respir Crit Care Med 2003;168(9):1095–1099. (Barnes-Jewish Hospital Foundation)</p>	<p>Prospective randomized, controlled trial</p>	<p>96 (ITT)</p>	<p>Age Mean = 36 yr</p> <p>Gender 18% male, 82% female</p> <p>Ethnicity 82% African American</p> <p>Education 62% 12th grade or lower</p>	<p>Moderate to severe airflow obstruction</p> <p>Asthma duration mean = 17 yr</p> <p>Age at onset of asthma, mean = 15 yr</p> <p>FEV₁ % pred., range 14–113, mean = 57</p> <p>FEV₁ mean = 1.56L</p> <p>All hospitalized in previous 12 months, mean = 2.4 times</p>
<p>C. ASTHMA SELF-MANAGEMENT EDUCATION IN EMERGENCY DEPARTMENT SETTINGS</p>				
<p>Baren et al. A randomized, controlled trial of a simple emergency department intervention to improve the rate of primary care follow-up for patients with acute asthma exacerbations. Ann Emerg Med 2001;38(2): 115–122. (American Lung Association, Southeastern Pennsylvania Chapter)</p>	<p>Randomized, controlled trial (urban university-based ED)</p>	<p>192 (192; ITT analysis)</p>	<p>Age 16–46 yr, mean = 30.8 yr</p> <p>Gender 28% male, 72% female</p> <p>Ethnicity 84% Black, 10% White, 4% Asian, 2% Hispanic</p> <p>Insurance Government-HMO, 40%; Government/military, 4%; HMO, 22%; Private, 22%; None, 12%</p>	<p>Acute asthma exacerbation</p> <p>Respiratory rate, 21.3 breaths/min</p> <p>O₂ saturation, mean = 96.8%</p> <p>PEFR, mean = 245 L/min</p> <p>Number of times inhaler used in past 24 hrs, mean = 4.8</p> <p>Inhaled steroids, 42%</p> <p>Oral steroids, 13%</p> <p>Symptom duration: < 6 hrs, 22%; 6–12 hrs, 10%; 12–24 hrs, 15%; > 24 hrs, 53%</p>
<p>Numata et al. Teaching time for metered-dose inhalers in the emergency setting. Chest 2002;122(2):498–504. (Fonds de la Recherche en Santé du Québec)</p>	<p>Descriptive report of data from 1 group in a randomized, open-label, parallel-group controlled trial (single acute-care day hospital)</p>	<p>61 (57) (Study sample included 119 with asthma and 68 with COPD. Only results of those who received the teaching protocol are reported. Table includes only those with asthma.)</p>	<p>Age ≥18 yr, median 46 yr</p> <p>Gender 65% male, 35% female</p> <p>Education Mean = 12.7 yr</p>	<p>Acute airflow obstruction</p> <p>FEV₁ % pred., mean = 63.5</p> <p>49% received 1 dose of bronchodilator</p> <p>Mean modified Borg scale score, = 3.7</p> <p>Mean = SpO₂ 96.1</p> <p>Previous MDI instruction, 65.6%</p> <p>Inhaled corticosteroid use during previous week, 73.8%</p> <p>Oral corticosteroid use during previous week, 23.0%</p>

Citation (Sponsor)	Study Design	Study Population		
		Study N (Number Evaluable)	Population Characteristics	Asthma Severity at Baseline (if reported)
Baren et al. Randomized controlled trial of emergency department interventions to improve primary care follow-up for patients with acute asthma. Chest 2006;129(2): 257–265.	Randomized controlled trial (9 EDs, majority were tertiary teaching hospitals serving poor urban populations)	384 (384; ITT analysis)	<p>Age 11–39 yr, mean = 25 yr</p> <p>Gender 45% male, 55% female</p> <p>Ethnicity 57% Black, 25% White, 15% Hispanic, 3% other</p> <p>Education 76% high school graduates</p> <p>Household income Estimated mean = \$28,700</p> <p>Insurance 27% Private, 38% Medicaid, 10% other public, 23% none</p>	<p>Asthma exacerbation; 75% presented with severe symptoms</p> <p>Duration of symptoms <24 hours, 41%</p> <p>Received steroid treatment in ED, 86%</p> <p>Discharged receiving prednisone</p> <p>25% admitted for asthma in past year</p> <p>During past 4 weeks: 84% inhaled beta₂-agonists, 34% inhaled corticosteroids</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Ford et al. Health outcomes among African American and Caucasian adults following a randomized trial of an asthma education program. Ethn Health 1997; 2(4):329–339. (Agency for Health Care Policy and Research)	Purpose/Objective: To conduct a race-specific re-analysis of a clinical trial of an asthma education program.				
	Intervention group (E): 3 small-group educational sessions on asthma led by a specially trained health care professional at either of the 2 ED sites. Those not attending were mailed the class materials. (n=79 African Americans, n=40 Caucasians)	Data collected through telephone interviews at baseline and 4, 8, and 12 months.			
	Control group (C): Usual care (n=84 African Americans, n=38 Caucasians)			E showed a decrease in number of ED visits vs. C (E: 5.0 to 2.7, C: 6.7 to 4.8 for African Americans; E: 4.7 to 1.5, C: 6.5 to 4.7 for Caucasian). There was no racial difference for E vs. C in number of ED visits (p=0.60), change in asthma knowledge and beliefs score (p=0.51), or limited activity days due to asthma (p=0.43) over the period of the study.	

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Levy et al. A randomized controlled evaluation of specialist nurse education following accident and emergency department attendance for acute asthma. <i>Respir Med</i> 2000; 94(9): 900–908.</p> <p>(National Asthma Campaign, UK; Allen and Hanbury's; Clement Clarke International; Astra Pharmaceuticals)</p>	<p>Purpose/Objective: To investigate whether patient education by hospital based specialist asthma nurses, using guided self-management plans based on nationally agreed guidelines, could improve patient recognition and self-treatment of asthma.</p>	<p>Interviews at baseline, 6 weeks, 3 months and 6 months</p>	<p>E vs. C had higher (mean 20.1 L/min, 95% CI 0.4 to 39.7) and less variable PEF ($p < 0.001$) at 6 months.</p>	<p>*Of those with mild attacks (E 14% & C 18%), E increased use of inhaled steroids vs. C (47% vs. 23%; OR 3.06, 95% CI 0.79 to 11.73) and use of rescue medication (89% vs. 82%; OR 1.89, 95% CI 0.31 to 11.68).</p> <p>Of those with severe attacks (E 34% & C 42%), E increased use of inhaled topical steroids vs. C (51% vs. 21%, OR 3.91, 95% CI 1.8 to 8.1, $p < 0.001$) and use of rescue medication (89% vs. 76%, OR 2.88, 95% CI 1.1 to 7.9), $p < 0.04$.</p> <p>E vs. C had lower ($p = 0.01$) and less variable ($p = 0.01$) symptom scores at 6 months.</p> <p>E had fewer days off work vs. C in the first 3 months (0.34 vs. 0.54, $p = 0.08$) but not in the second 3 months (0.25 vs. 0.30).</p> <p>E vs. C had fewer routine consultations with the doctor ($p < 0.05$) and nurse ($p < 0.03$) with no difference in uncontrolled episodes and hospital visits.</p>	<p>Quality of life scores improved in both groups with no difference between groups.</p>
	<p>Intervention group (E): 1-hour consultation with nurse followed by 2 more consultations lasting 30 minutes at 6-week intervals. Patients taught to step up medication based on PEF or symptoms. (n=103, n=79 attended at least 1 consultation, n=81 completed interviews)</p> <p>Control group (C): Usual care (n=108, n=94 completed interviews)</p>				

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Klein et al. Benefit from the inclusion of self-treatment guidelines to a self-management programme for adults with asthma. Eur Respir J 2001; 17(3):386–394.</p> <p>(Netherlands Asthma Foundation; GlaxoWellcome; the “Stichting Astmabestrijding”; Amicon Health Care Insurance Fund)</p>	<p>Purpose/Objective: To assess the long-term effects and safety of adding self-treatment guidelines to a self-management program for adults with moderate to severe asthma who were well controlled and already on a relatively high dose of inhaled corticosteroids.</p>		<p>Decrease in PEF variability over 24 months was 32% in E and 29% in C. Difference between groups not significant.</p> <p>PC₂₀ histamine increased significantly in both groups over 12 months (2.54 mg/mL in E; 2.33 mg/mL in C). Difference between groups not significant.</p> <p>C showed a significant increase in FEV₁ over E (3% vs. 2%).</p>	<p>Frequency of exacerbations decreased by 2% in E and increased by 5% in C (diff 7%, 95% CI –2 to 16).</p> <p>Difference in total number of outpatient visits in both groups decreased (E by 25%, C by 18%) with no difference between groups.</p> <p>No difference between groups in percentage of symptom free days and nights.</p>	<p>Quality of life for both groups increased by 7% over the 12 months with no difference between groups in overall score or domain scores.</p> <p>After 12 months, percentage who perceived better control of asthma was 90.7% for E and 74.6% for C (p<0.001). Perceived self-confidence was 3.76 for E vs. 3.41 for C (p=0.018).</p>
	<p>Self-treatment group (E): Instructions about self-treatment of exacerbations: told to measure PEF weekly, both on a fixed day and any time asthma symptoms worsened, and given a 4-zone self-treatment plan. (n=123; n=? completers)</p> <p>Control group (C): Usual care (n=122; n=? completers)</p>	<p>2-week diary data and spirometry collected at 4, 8, 12, 18, and 24 months after start of educational program</p> <p>All subjects were educated by a specially trained asthma nurse in 3 consecutive weekly sessions each lasting 90 minutes. Self-treatment instruction given during 3rd session to E.</p>			
<p>Neri et al. Short and long-term evaluation of two structured self management programmes on asthma. Monaldi Arch Chest Dis 2001;56(3): 208–210.</p>	<p>Purpose/Objective: To evaluate the efficacy of 2 structured self-management programs on asthma at the end of interventions (short-term efficacy) and after 3 years (long-term efficacy).</p>		<p>*No difference between groups in FEV₁ at 1 year and at 3 years (p>0.05).</p>		<p>*No difference between groups in knowledge at 1 year and at 2 years (p>0.05).</p> <p>*No difference between groups in number of correct steps performed during the inhalation procedure at 1 year and at 3 years (p>0.05).</p>
	<p>Intervention group (E): “Asthma school” program consisting of 6 lessons of 1 hour each based on content in line with the International Asthma Guidelines (n=20)</p> <p>Comparison group (C): “Basic” educational program (n=20)</p>	<p>1-year and 3-year assessments</p>			

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Van der Palen et al. Behavioural effect of self-treatment guidelines in a self-management program for adults with asthma. Patient Educ Couns 2001; 43(2):161–169.</p> <p>(The Netherlands Asthma Foundation; the “Stichting Astmabestrijding Nederland”; GlaxoWellcome)</p>	<p>Purpose/Objective: To assess whether including self-treatment guidelines (action plans) in a self-management program for adult asthmatics leads to greater behavioural changes than a similar program without these guidelines.</p>	<p>2-week diary 1 year after intervention</p>			<p>No difference between E and C in change in generalized self-efficacy (3.5% vs. 1.6%), asthma-specific self-efficacy expectancies (14% vs. 13%), and outcome expectancies (4% for both E and C) over 1 year.</p> <p>No change in intention toward self-management in either group.</p> <p>No differences between E and C in change in asthma-specific knowledge (20.3% for E and C combined).</p>
	<p>Intervention group (E): Self-management training and education by a specialized asthma nurse in three 90-minute small group sessions, instruction in correct use of inhaled medication and a peak flow meter, and instructions about the self-treatment of exacerbations. (n=123; 119 completers)</p> <p>Comparison group (C) Self-management training and education by a specialized asthma nurse in three 90-minute small group sessions and instruction in correct use of inhaled medication and a peak flow meter. No instruction about the self-treatment of exacerbations. (n=122; n=119 completers)</p>				

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Couturaud et al. Education and self-management: a one-year randomized trial in stable adult asthmatic patients. J Asthma 2002; 38(6): 493–500. (Projets Hospitaliers de Recherche Clinique 1995)	<p>Purpose/Objective: To assess the effects of an educational program in asthmatic patients following treatment readjustment.</p> <p>Intervention group (E): Personal education program consisting of 5 sessions covering pathophysiology of asthma, role of medication and side effects, asthma triggers and their avoidance, detection of an asthma flare up, and self-management plan based on symptoms and peak-flow monitoring (n=36; 26 completers)</p> <p>Control group (C): No personal education program (n=36; 28 completers)</p>	<p>30–60 minutes of education at each of 5 sessions</p> <p>2-week run in period to optimize asthma control with a maximum of 3 run in periods</p> <p>12-month followup</p>	Control and Education groups comparable in FEV ₁ (84% vs. 85%, p>0.05).	<p>*No difference in symptom-free days (89% for E and 88% for C).</p> <p>No difference in days off work (4% for E, 8% for C) or unscheduled visits (1 vs. 0.6 for E vs. C).</p> <p>Percentage of days of oral steroid intake higher in E vs. C (6% vs. 3%, p=0.01).</p> <p>Among E, those who complied with action plan (n=5) had higher percentage of symptom-free days (97% vs. 87%, p=0.009).</p>	Asthma Quality of Life scores did not differ between E and C.

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Cowie et al. Asthma in adolescents: a randomized, controlled trial of an asthma program for adolescents and young adults with severe asthma. Can Respir J 2002;9(4):253–259. (Alberta, Canada, Lung Association)	Purpose/Objective: To determine the impact of an age-specific asthma program on asthma control, particularly on exacerbations of asthma requiring ED treatment, and on the quality of life of adolescents with asthma.				At 3 months, 12.5% of E vs. 34.0% of C had visited an ED for asthma treatment (p=0.03). At 6 months, there were no differences between E and C in disease control and management.
	Intervention group (E): Young Adult Asthma Program was staffed by asthma educators, respiratory therapists, and a respiratory physician or pediatrician. Subject assessed individually and seen by a physician trained in asthma management for 90–120 minutes. Report was sent to regular family physician. Subject given appointment for followup visit with full team. (n=?; n=29 completers)	Asthma and quality of life questionnaires mailed at 3 months and at 6 months.			
	Control group (C): Usual care (n=?; n=29 completers)				

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Gibson et al. Self-management education and regular practitioner review for adults with asthma (Review). Cochrane Database Syst Rev 2000(2) or 2003(1): CD001117. (NSW Health Cooperative Research Centre for Asthma, Australia)	Purpose/Objective: To evaluate the literature supporting Step 5 of the Australian Asthma Management Plan, "Educate and Review Regularly," to determine whether health outcomes are influenced by asthma education and self-management programs.		No significant effect of education on FEV ₁ (10 studies) or PEF (16 studies)	Self-management education reduced hospitalizations (RR 0.64, 95% CI 0.50 to 0.82; 8 studies), emergency room visits (RR 0.82, 95% CI 0.73 to 0.94; 20 studies), unscheduled visits to the doctor (RR 0.68, 94% CI 0.56 to 0.81; 3 studies), and days off work or school (RR 0.79, 95% CI 0.67 to 0.93; 16 studies).	Self-management education increased quality of life (standard mean difference 0.29, 95% CI 0.11 to 0.47; 3 studies).
	Intervention group (E): Self-management education included optimal self-management (15 studies), self-monitoring and regular review (7 studies), self-monitoring only (10 studies), regular review only (1 study), and written action plan but not optimal self-management (2 studies)	Control group (C): Usual care			

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Gibson et al. Limited (information only) patient education programs for adults with asthma (Review. Cochrane Database Syst Rev 2002;(2): CD001005. (Cooperative Research Centre for Asthma, Australia; Garfield Weston Foundation UK)	<p>Purpose/Objective: To assess the effects of limited (i.e., information only) asthma education on health outcomes in adults with asthma.</p> <p>Intervention (E): Asthma education delivered to individual or group that included information about pathophysiology of asthma, management of trigger factors, and action and side effects of medication</p> <p>Control (C): Usual care, which may or may not have involved a degree of education, waiting list control, or lower-intensity educational intervention</p>		E did not affect FEV ₁ (weighted mean difference 0.20, 95% CI -0.35, 0.75; 1 study).	E did not reduce hospitalization for asthma (weighted mean difference -0.03 hospitalizations/per person/year, 95% CI -0.09, 0.03; 3 studies) or doctor visits (weighted mean difference 0.22 visits/person/year, 95% CI -0.09, 0.52; 5 studies). No difference in medication use (n=3 studies). No reduction in days lost from normal activity (n=3 studies).	
Marabini et al. Short-term effectiveness of an asthma educational program: results of a randomized controlled trial. Respir Med 2002; 96(12):993-998. (National Research Council, Italy)	<p>Purpose/Objective: To assess the effectiveness of an educational program and its impact on quality of life.</p> <p>Intervention group (E): Usual treatment plus educational program based on "Teach Your Patients about Asthma: A Clinician Guide." Education delivered by physician. (n=37)</p> <p>Control group (C): Usual care (n=40)</p>	Three 2-hour sessions Followup data collected at 3 months	No significant improvement in FEV ₁ or PEF in either group. When stratified for asthma severity, FEV ₁ was higher in E vs. C (81.2 vs. 72.5, p<0.05) after adjusting for baseline.	No significant improvement in morbidity in either group.	At 3 months, E vs. C had higher score in quality of life overall (6.0 vs. 5.6, p<0.05) and in environment (6.6 vs. 6.0, p<0.005) and symptom (6.2 vs. 5.7, p<0.005) domains. Among those with moderate-to-severe asthma, E improved in overall QOL score (5.6 to 6.0, p<0.05), symptom score (5.7 to 6.2, p<0.01), and activities (5.1 to 5.4, p<0.05) domains with no change in scores of C.

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Perneger et al. Effect of patient education on self-management skills and health status in patients with asthma a randomized trial. Am J Med 2002; 113(1):7-14. (Swiss National Science Foundation, Berne, Switzerland)</p>	<p>Purpose/Objective: To assess the effectiveness of a newly established education program for adults with asthma designed to improve patients' health and functional status.</p> <p>Intervention group (E): Three 75-minute educational group sessions, 1 week apart, conducted by 2 respiratory physicians and a physiotherapist. Purpose was to design individual self-management plans. Copies of plans were sent to the patients' physicians. (n=66; n=57 completers)</p> <p>Control group (C): Delayed education scheduled after followup assessment at 6 months. (n=65; n=58 completers)</p>	<p>Assessed at 6 months postintervention.</p>	<p>No difference between E and C in improvements in pulmonary function</p>		<p>2 of 9 improvements in self-management favored E: "correct inhalation technique" (OR 2.4, 95% CI 1.0 to 5.7, p=0.048) and "knows peak flow reading that requires calling physician" (OR 3.1, 95% CI 1.4 to 6.7, p=0.004). No difference between E and C in functional outcomes or use of health care in past 6 months.</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Powell et al. Options for self-management education for adults with asthma (Review). Cochrane Database Syst Rev 2003;(1) CD004107. (Cooperative Research Centre for Asthma, Australia)	Purpose/Objective: To compare the differing components of self-management interventions in asthma.		Comparison 2: No difference in PEF (6 studies). Comparison 3: Regular review was associated with improvement in mean percent predicted FEV ₁ as compared with omission of regular review (1 study).	Comparison 1: No difference in hospitalizations for asthma (4 studies), emergency room visits (1 study), unscheduled doctor visits (studies), and nocturnal asthma (3 studies). Comparison 3: Omission of regular review was associated with more health center visits and 2.5 more sickness days (1 study).	
	Three primary comparisons: (1) Optimal Self-Management vs. Regular Medical Review (6 studies) (2) Peak Expiratory Flow Based vs. Symptom Based Self-Management (6 studies) (3) Optimal Self-Management vs. Modified Optimal Self-Management (3 studies)	6-month followup visit to education program Health care utilization and absenteeism during 6 months before and 6 months after initial visit compared.			
	3 levels of education based on patient asthma severity and initial needs assessment. Topics included trigger avoidance, environmental control, the role of medications, delivery systems, action plans, and self-monitoring skills. Assessment and education took approximately 2 hours.				

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Janson et al. Effects of individual self-management education on clinical, biological, and adherence outcomes in asthma. Am J Med 2003;115(8): 620–626. (National Institutes of Health)</p>	<p>Purpose/Objective: To assess the effects of individual self-management education on adherence to anti-inflammatory medication, biological markers of airways inflammation, and clinical outcomes.</p> <p>Intervention group (E): 30-minute individual session of asthma education consisting of components recommended in National Institutes of Health guidelines: basic facts about asthma, role of airway inflammation and bronchospasm, roles and actions of anti-inflammatory and quick relief medications, skill for correct inhalation and for peak flow measurement. Reinforcement by study coordinator at next 2 biweekly visits. (n=33)</p> <p>Control group (C): Monitoring only (n=32)</p>	<p>6 weeks with 5 study visits after 1-week run in.</p>	<p>*No difference between groups in change in FEV₁ % pred. (p=0.09) or morning peak flow (p=0.23).</p>	<p>*Adherence increased in E from 70% to 91% vs. decrease from 65% to 62% in C (p=0.01). Perceived control of asthma improved 14% in E vs. 5% in C (p=0.04). No difference between E and C in change in use of beta-agonist (p=0.21).</p>	<p>No difference in increase in quality of life (37% for E and 21% for C, p=0.06).</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Thoonen et al. Self-management of asthma in general practice, asthma control and quality of life: a randomized controlled trial. Thorax 2003; 58(1):30-36.</p> <p>(The Netherlands Organization for Scientific Research; ASTRAZeneca Pharmaceutica BV)</p>	<p>Purpose/Objective: To determine if guided self-management can provide a safe treatment strategy for asthmatic patients in general practice.</p> <p>Intervention group (E): 4 individual training sessions during a period of 3 months consisting of tailored education and instructions on how to use a personalized written self-treatment plan. After training sessions, biannual control visits were recommended over a followup period of 21 months. (n=95; n=86 completers)</p> <p>Control group (C): Usual care according to guidelines of the Dutch College of Family Physicians, which recommends followup visits every 3-6 months. 1 visit was scheduled at start to instruct patients on the use and dosage of inhaled steroids (budesonide 200 mcg Turbuhaler). (n=98; n=85 completers)</p>	<p>Assessment every 6 months for 2 years</p>	<p>No difference between E and C in estimated rate of decline in FEV₁ reversibility and PC₂₀ histamine.</p>	<p>No difference between E and C in median dose of short-acting beta₂ bronchodilators (p=0.71) or ipratropium (p=0.61), but a higher number of courses of oral prednisolone in E vs. C (p=0.015).</p>	<p>*Increased number of mean percentage of successfully treated weeks per patient in E vs. C (78% vs. 72%, p=0.003).</p> <p>*Mean number of limited activity days was lower in E vs. C (1.2 vs. 3.9; p<0.05).</p> <p>No difference between E and C in number of diagnosed exacerbations.</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Hopman et al. Outcomes of asthma education: results of a multisite evaluation. Can Respir J 2004; 11(4):291–297.	Purpose/Objective: To compare the effectiveness of a standardized asthma education program on health care utilization, absenteeism, amount of leisure time missed, and health-related quality of life in a variety of settings.			*Significant improvement ($p < 0.001$) in regular physician visits (3.13 vs. 1.96), unscheduled physician visits (2.71 vs. 0.51), number of hospital admissions (0.46 vs. 0.13), number of days in hospital (1.72 vs. 0.92), number of emergency room visits (0.96 vs. 0.27), missed work/school days (5.04 vs. 3.85), and missed leisure time days (2.14 vs. 1.61). Significant improvement in unscheduled specialist visits (0.14 vs. 0.06, $p < 0.05$). No significant difference in proportion taking inhaled corticosteroid (90.5% vs. 87.2%). Use of long-acting beta ₂ -agonist increased from 11.5% to 20.6% ($p = 0.02$) and use of oral corticosteroid decreased from 17.9% to 6.2% ($p = 0.02$).	6 months after asthma education there were significant improvements in all health-related quality of life domains (SF-36) except mental health. Change in physical function, role physical, role emotional, bodily pain, and vitality exceeded clinically meaningful change of 5 points.
	3 levels of education based on patient asthma severity and initial needs assessment. Topics included trigger avoidance, environmental control, the role of medications, delivery systems, action plans, and self-monitoring skills. Assessment and education took approximately 2 hours.	6-month followup visit to education program Health care utilization and absenteeism during 6 months before and 6 months after initial visit compared.			
Magar et al. Assessment of a therapeutic education programme for asthma patients: “un soufflé nouveau.” Patient Educ Couns 2005;58(1):41–46. (GlaxoSmithKline)	Purpose/Objective: To assess the effects of a new asthma self-management education program entitled “un soufflé nouveau.”				Number of symptom-free days increased over time in E, but not C (3.5 vs. -0.22 at 6 months; 3.53 vs. -0.26 at 12 months, $p = 0.03$). Number of nighttime awakenings decreased in E vs. C (-2.11 vs. -0.48 at 6 months; 0.1 vs. -0.17 at 12 months, $p = 0.04$). Use of oral corticosteroids decreased in both groups with a gain of 8.8 days in E vs. 1.8 days in C ($p = 0.03$). Monthly use of fast-acting beta ₂ -agonists decreased in E from 10.6 at 6 months to 1.3 at 12 months with no change in C ($p = 0.03$).
	Intervention group (E): Education program of initial 30–60 minute appointment followed by 2 group sessions of 2.5 hours each at an interval of 15 days, each addressed selected teaching objectives. ($n = 127$; $n = 104$ completers)	Both groups assessed at 6 months and 12 months.			
	Control group (C): Information in form of printed document. ($n = 111$; $n = 89$ completers)				

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Urek et al. Effect of educational programs on asthma control and quality of life in adult asthma patients. Patient Educ Couns 2005;58(1):47-54.	<p>Purpose/Objective: To investigate the effectiveness of different forms of medical education in improving the control of asthma and quality of life in patients with moderate persistent asthma.</p> <p>Group 1 (AS) Asthma school: Three 4-hours sessions of group education consisting of lectures and workshops. (n=20; n=20 completers)</p> <p>Group 2 (IVI) Individual verbal instruction in three 1-hour sessions. (n=20; n=20 completers)</p> <p>Group 3 (B) Written information in an asthma booklet. (n=20; n=20 completers)</p>	<p>12 weeks</p> <p>All patients regularly treated with inhaled corticosteroids and allowed to use salbutamol as rescue medication as needed. No change in medication allowed during the study.</p>	<p>Using repeated measures and adjusting for gender, age, and baseline values, improvement was greater for IVI than for B in morning PEFR (p<0.05) and evening PEFR (p<0.05).</p>		<p>All groups improved in daily average number of day and nighttime symptoms (p<0.001) with no difference between groups (p=0.74 for daytime; p=0.56 for nighttime) using repeated measures and adjusting for gender, age, and baseline values.</p> <p>All groups decreased significantly in use of rescue medication (p<0.05) with no difference between groups (p=0.61 for day usage; p=0.87 for night usage) using repeated measures and adjusting for gender, age, and baseline values.</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
George et al. A comprehensive educational program improves clinical outcome measures in inner-city patients with asthma. Arch Intern Med 1999; 159(15): 1710–1716. (NIH; Health Management Alternatives, Abbott Laboratories; Mary Rockefeller Foundation; Merck & Co, West Point, PA; Schering-Plough/Key Pharmaceuticals, Kenilworth, NJ)	<p>Purpose/Objective: To examine whether a comprehensive in-patient educational program by an asthma nurse educator improved outpatient follow-up to an asthma outpatient program and whether outpatient follow-up then decreased subsequent hospital utilization in a cohort of indigent, inner-city patients with asthma.</p> <p>Intervention group (E): Educational program that included inpatient asthma instruction, bedside spirometry, assistance with discharge planning, a post discharge telephone call, and scheduled follow-up in an outpatient asthma care program (n=44; n=30 with 6 month data)</p> <p>Control group (C): Routine care (n=33; n=20 with 6 month data)</p>	Assessment at 6 months after enrollment			<p>Mean length of hospital stay did not differ between groups (2.1 for E vs. 2.7 for C, p=0.12).</p> <p>*60% of E vs. 27% of C returned for initial follow-up visit (p=0.01).</p> <p>Patients in E vs. C had reduced need for ED visits (p=0.04) and number of hospitalizations (p=0.04) in 6 months after enrollment as compared to 6 months prior to enrollment.</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Morice & Wrench. The role of the asthma nurse in treatment compliance and self-management following hospital admission. <i>Respir Med</i> 2001;95(11): 851–856. (Astra Pharmaceuticals)</p>	<p>Purpose/Objective: To determine primarily whether nurse intervention could change behavior in adult asthmatics and whether this was reflected in the number of emergency visits/call-outs to General Practitioners and re-admission to hospital with asthma-related programs following discharge.</p> <p>Intervention group (E): Education program from the asthma nurse over a minimum of 2 separate sessions, lasting on average 30 minutes each, carried out on an individual basis (n=40; n=39 at 6 weeks; n=35 at 6 months)</p> <p>Control group (C): Information in form of printed document (n=40; n=31 at 6 weeks; n=30 at 6 months)</p>	<p>Data collection at 6 weeks and 6 months; General Practitioners contacted at 4 months.</p>			<p>Higher proportion presented to hospital in E vs. C (91% vs. 62.5%) over the 18 months of the study.</p> <p>No difference in combined endpoint of readmissions plus emergency call-outs over the 18 months of the study.</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Osman et al. A randomized trial of self-management planning for adult patients admitted to hospital with acute asthma. Thorax 2002; 57(10): 869–874. (NHS R&D Programme in Delivery of Care in Asthma managed by the National Asthma Campaign, UK)	<p>Purpose/Objective: To test whether the self-management program offered in the hospital to adults admitted with acute asthma was effective in reducing readmission and morbidity after discharge and whether it was acceptable to patients as part of their hospital management.</p> <p>Intervention group (E): Self-management program of 2 visits of about 30 minutes each by a trained respiratory nurse that included a symptom-and peak-flow-based self-management plan. (n=135; n=108 at 1 month; n=131 at 12 months)</p> <p>Control group (C): Standard care as directed by patient's physician (n=145; n=118 at 1 month; n=140 at 12 months)</p>	Patients surveyed at 4 weeks after discharge; hospital records reviewed at 12 months			<p>*At 12 months E less likely than C to be readmitted (17% vs. 27%, OR 0.5, 95% CI 0.3 to 1.0, p=0.04).</p> <p>At discharge, E more likely than C to be prescribed inhaled steroids (99% vs. 92%, p=0.03) and to have hospital follow up (98% vs. 84%, p<0.01).</p> <p>At 1 month, E more likely than C to report no daytime wheeze (OR 2.6, 95% CI 1.2 to 5.3) and no night disturbance (OR 2.0, 95% CI 1.2 to 3.5) and more likely to have written management plan (93% vs. 30%, OR 28.0, 95% CI 12.3 to 63.8).</p> <p>E vs. C more satisfied with hospital explanation of asthma (100% vs. 76%, p<0.001).</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Castro et al. Asthma intervention program prevents readmissions in high healthcare users. Am J Respir Crit Care Med 2003;168(9): 1095–1099. (Barnes-Jewish Hospital Foundation)</p>	<p>Purpose/Objective: To study the effect of a limited nurse-focused intervention on rate of admission, lost work/school days, asthma-specific quality of life, and overall cost of health care.</p> <p>Intervention group (E): Multifaceted approach by asthma nurse specialist including “Asthma Care” flow sheet while in hospital, asthma education, psychosocial support, individualized self-management plan, facilitation of discharge planning, and outpatient followup. (n=50)</p> <p>Control group (C): Usual care (n=46)</p>	<p>Assessment at 6 and 12 months; costs during 180 days.</p>		<p>*60% reduction in readmissions in E vs. C (31 vs. 71, p=0.04). Multiple readmissions more frequent in C vs. E (15 vs. 7, p=0.03). Adjusting for other variables, C was 3.6 times more likely to be readmitted <u>2</u> times or more (95% CI 1.2–10.9).</p> <p>Time to first readmission was not different between E and C.</p> <p>69% reduction in hospital days in E vs. C (p=0.04).</p> <p>No difference in number of health care provider visits.</p> <p>Fewer lost work/school days for E vs. C (246 vs. 1040).</p> <p>Savings of \$6462/patient attributable to the intervention (p=0.03).</p>	<p>No difference in improvement of quality of life between 2 groups (1.2 for E vs. 1.4 for C, p=0.55).</p>

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Baren et al. A randomized, controlled trial of a simple emergency department intervention to improve the rate of primary care follow-up for patients with acute asthma exacerbations. Ann Emerg Med 2001;38(2): 115–122. (American Lung Association, Southeastern Pennsylvania Chapter)</p>	<p>Purpose/Objective: To determine whether a simple, inexpensive, 3-part ED intervention would increase the likelihood of PCP followup for patients treated and discharged from the ED for an acute asthma exacerbation.</p> <p>Intervention group (E): Self-contained “fanny pack” of materials designed to enhance the likelihood of scheduling timely followup: 5-day course of oral prednisone, 2 taxicab vouchers, an asthma information card, written instructions for use of medications and vouchers, and telephone reminder within 48 hours of ED release to make appointment with PCP. (n=98; n=95 completers)</p> <p>Control group (C): ED discharge instructions and medication prescriptions at discretion of treating physician. (n=94; n=83 completers)</p>	<p>Contact at 4 weeks by investigator blinded to study group assignment.</p>		<p>*46.3% of E and 28.9% of C followed up with PCP (RR 1.6, 95% CI 1.1 to 2.4).</p> <p>Characteristics associated with followup were older age, prior relationship with PCP (RR 4.4, 95% CI 1.9 to 10.2), black race (RR 0.54, 95% CI 0.37 to 0.79), and lack of health insurance (RR 0.33, 95% CI 0.11 to 0.96).</p> <p>Adjusting for variables associated with PCP followup, E was associated with increased likelihood of followup (adj. OR 3.0, 95% CI 1.5 to 6.3).</p>	

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
<p>Numata et al. Teaching time for metered-dose inhalers in the emergency setting. Chest 2002;122(2): 498–504. (Fonds de la Recherche en Santé du Québec)</p>	<p>Purpose/Objective: To measure the teaching time that was necessary for successful MDI/DED use at acute-care facility based on demonstrated mastery according to a predefined protocol.</p> <p>Intervention group (E): MDI/DED with standardized one-on-one teaching from 1 of 3 RNs based on a protocol of nurse demonstration, patient return demonstration, and teaching about care and maintenance of the device. (n=62; 57 completers)</p> <p>Control group (C): Wet nebulizer (n=57; n=56 completers) (Results for C not reported)</p>	<p>Teaching directly observed and timed. Treating physicians selected the bronchodilator(s) used and frequency of their administration; standard doses were used.</p>		<p>*Median total teaching time was 6.2 min (IQR 4.3 to 9.2 min). Among those with no previous MDI instruction, median teaching time was 8.4 min (IQR 5.9 to 11.3). 5% of patients required inpatient hospitalization. Most patients expressed satisfaction with MDI/DED teaching and treatment.</p>	

Citation/Sponsor	Study Characteristics		Findings		
	Treatment	Duration of Active Treatment; Duration of Postintervention/ Off-Treatment Followup	Lung Function	Morbidity	Knowledge/Quality of Life
Baren et al. Randomized controlled trial of emergency department interventions to improve primary care follow-up for patients with acute asthma. Chest 2006; 129(2):257–265.	Purpose/Objective: To compare the effect of 2 ED interventions on the rates of primary care follow-up for patients receiving either telephone reminders to make primary care appointments vs. those receiving a telephone contact to give them an actual appointment.			<p>*Patients in E2 were most likely to have a followup visit (E2, 73% vs. C, 49% and E1, 49%, p<0.001).</p> <p>Adjusting for age, race, gender, prior PCP, and insurance status, intervention for E2 remained significant (OR 2.8, 95% CI 1.5 to 5.1).</p> <p>Among patients with 30-day follow-up who did not have a PCP at enrollment, those in E2 were more likely to visit a PCP (45% of E1 vs. 30% of E1 and 19% of C, p=0.06).</p> <p>No differences in ED, hospitalizations, quality of life, or inhaled corticosteroid use at 12 months after index ED visit.</p>	
	<p>Intervention group 1 (E1): 5-day course of prednisone, 2 transportation vouchers for travel to and from PCP, general information sheet, telephone contact 2 days after visit to advise patient to make followup appointment (n=126; n=110 at 30 days; n=82 at 12 months)</p> <p>Intervention group 2 (E2): 5-day course of prednisone, 2 transportation vouchers for travel to and from PCP, general information sheet, telephone contact 2 days after visit to give a follow-up appointment if patient had not made one (n=132; n=106 at 30 days; n=87 at 12 months)</p> <p>Control group (C): Usual discharge care—not standardized (n=126; n=98 at 30 days; n=78 at 12 months)</p>	Followup telephone interview 2 days, 30 days, and 12 months following ED visit; PCPs contacted at 30 days to verify followup is obtained.			