


National Asthma Education and
Prevention Program

MANAGEMENT OF ASTHMA EXACERBATIONS: *School Treatment*

Suggested Emergency Nursing Protocol for Students with
Asthma Symptoms Who Don't Have a Personal Asthma Action Plan

A student with asthma symptoms should be placed in an area where he/she can be closely observed. Never send a student to the health room alone or leave a student alone. Limit moving a student who is in severe distress. Go to the student instead.

See list of **Possible Observations/Symptoms** on back.



ED: emergency department
PEF: peak expiratory flow

MDI: metered-dose inhaler
SABA: short-acting beta2-agonist (quick-relief inhaler)

AUGUST 2011

Possible Observations/Symptoms (May include one or more of the following):

- » Coughing, wheezing, noisy breathing, whistling in the chest.
- » Difficulty or discomfort when breathing, tightness in chest, shortness of breath, chest pain, breathing hard and/or fast.
- » Nasal flaring (nostril opens wide to get in more air).
- » Can only speak in short phrases or not able to speak.
- » Blueness around the lips or fingernails.

***Risk Factors for Death from Asthma**

Asthma history

- » Previous severe exacerbation (e.g., intubation or ICU admission for asthma).
- » Two or more hospitalizations for asthma in the past year.
- » Three or more ED visits for asthma in the past year.
- » Hospitalization or ED visit for asthma in the past month.
- » Using >2 canisters of SABA per month.
- » Difficulty perceiving asthma symptoms or severity of exacerbations.
- » Other risk factors: lack of a written asthma action plan, sensitivity to Alternaria.

Social history

- » Low socioeconomic status or inner-city residence.
- » Illicit drug use.
- » Major psychosocial problems.

Comorbidities

- » Cardiovascular disease.
- » Other chronic lung disease.
- » Chronic psychiatric disease.

**** Consider administering epinephrine** if the student is unable to use SABA because respiratory distress or agitation prevents adequate inhalation from the SABA inhaler device and nebulized albuterol is not available and the exacerbation is **life-threatening**. Administer epinephrine auto-injector in lateral thigh as per local or state epinephrine protocol. Epinephrine is NOT first line treatment for asthma. Albuterol is the treatment of choice. Administration of epinephrine should be rare and is intended to prevent a death at school from a severe asthma attack. Most school nurses will never need to administer epinephrine.