

# Medicine & Science IN Sports & Exercise

The Official Journal of the American College of Sports Medicine  
www.acsm-msse.org

**. . . Published ahead of Print**

## **Novel Strategies for Sedentary Behavior Research**

Dori E. Rosenberg<sup>1</sup>, I-Min Lee<sup>2</sup>, Deborah Rohm Young<sup>3</sup>, Thomas R. Prohaska<sup>4</sup>,  
Neville Owen<sup>5</sup>, and David M. Buchner<sup>6</sup>

<sup>1</sup>Group Health Research Institute, Seattle, WA; <sup>2</sup>Harvard Medical School, Boston, MA;  
<sup>3</sup>Kaiser Permanente Southern California, Pasadena, CA; <sup>4</sup>George Mason University,  
Fairfax, VA; <sup>5</sup>Baker IDI Heart and Diabetes Institute, Melbourne, Australia; <sup>6</sup>University  
of Illinois Urbana-Champaign, Champaign, IL

Accepted for Publication: 28 July 2014

**Medicine & Science in Sports & Exercise**® **Published ahead of Print** contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Copyright © 2014 American College of Sports Medicine

## Novel Strategies for Sedentary Behavior Research

Dori E. Rosenberg<sup>1</sup>, I-Min Lee<sup>2</sup>, Deborah Rohm Young<sup>3</sup>, Thomas R. Prohaska<sup>4</sup>,  
Neville Owen<sup>5</sup>, and David M. Buchner<sup>6</sup>

<sup>1</sup>Group Health Research Institute, Seattle, WA

<sup>2</sup>Harvard Medical School, Boston, MA

<sup>3</sup>Kaiser Permanente Southern California, Pasadena, CA

<sup>4</sup>George Mason University, Fairfax, VA

<sup>5</sup>Baker IDI Heart and Diabetes Institute, Melbourne, Australia

<sup>6</sup>University of Illinois Urbana-Champaign, Champaign, IL

**Corresponding author:** David Buchner, MD, MPH, FACSM  
Department of Kinesiology and Community Health, MC-588  
University of Illinois at Urbana-Champaign  
Phone: 217-244-1510 Email: dbuchner@illinois.edu

**Running Title:** Strategies for Sedentary Behavior Research

**Conflict of Interest:** None

## **Acknowledgements**

The workshop was sponsored by the National Heart, Lung and Blood Institute, the National Institute on Aging and the Office of Disease Prevention of the National Institute of Health.

The views expressed in this document reflect the collective ideas and opinions of the authors and does not necessarily represent the official views of the National Institutes of Health , or the U.S. Department of Health and Human Services and the other workshop participants. Support for this work provided in part by a Shahid & Ann Carlson Khan Professorship (Dr. Buchner), grant CA154647 from the US National Institutes of Health (Dr. Lee, by National Health and Medical Research Council of Australia Program Grant (#569940) and Research Fellowship funding (#1003960) and by Victorian Government OIS funding (Dr. Owen). Authors were provided with an honorarium from the National Institutes of Health for their participation in the workshop. We wish to thank the discussants for this workshop: Dr. Stephen P. Fortmann, Kaiser Permanente Center for Health Research, Dr. James F. Sallis, University of California San Diego, Dr. William Haskell, Stanford University School of Medicine, and Dr. Steven N. Blair, University of South Carolina. The results of the present study do not constitute endorsement by ACSM.

## **Abstract**

**Purpose:** This paper reports on the ‘Novel Strategies for Sedentary Behavior Research’ session of the *Sedentary Behavior: Identifying Research Priorities* workshop.

**Methods:** The purposes of this session of the workshop were to propose strategies for accomplishing a research agenda in dealing with sedentary behavior and to consider research priorities for people at high risk for excess sedentary behavior.

**Results and conclusions:** The four major recommendations from this workshop were: (1) Add repeated, objective measures of physical activity and sedentary behavior to existing cohort studies and standardize approaches to measurement and analysis. Epidemiologic studies will be the most efficient design for addressing some research questions. (2) To increase research efficiency, consider the advantages of a network of connected research studies and health systems. Advantages include access to existing data in electronic health records. (3) In intervention research, carefully select a variety of high-risk study populations and pre-plan collaboration among studies. This strategy can efficiently address the breadth of issues in sedentary behavior research. (4) In intervention research, include comparative effectiveness designs and pure environmental interventions. This strategy facilitates and enhances translation of interventions into practice.

**Key words:** physical activity, research priorities, sitting time, consensus

This paper reports on the proceedings of the last of four sessions as part of a joint workshop sponsored and organized by the National Heart, Lung, and Blood Institute and National Institute on Aging entitled *Sedentary Behavior: Identifying Research Priorities*. Presentations and subsequent discussions focused specifically on epidemiologic approaches, leveraging opportunities for research in health systems, translation of research findings, and international perspectives on sedentary behavior research strategies. The sessions also focused on promoting an efficient and inclusive research agenda. We present the key recommendations from the panel along with a brief rationale for each, which stemmed from the presentations and resulting discussion. We also include limited content that was not specifically discussed within the workshop but that the authors believe are important.

**Recommendation 1:** *To enhance the efficiency of observational research and facilitate pooling of data from cohort studies, in addition to self-report measures of sedentary behaviors, add repeated, objective measures of physical activity and sedentary behavior to existing cohort studies and standardize approaches to measurement and analysis.*

**Rationale:** As mentioned in the first workshop, “Definition, Measurement and Health Risks Associated with Sedentary Behavior”, a gap in the literature is the limited availability of datasets with objectively measured sedentary behavior (5). For the past decade, most research reports have relied on self-report assessments of sedentary behavior and several large cohorts include self-reports of sedentary behavior (15, 21, 36, 37). However, self-reported measures of sedentary behavior often have small associations with accelerometer measured sedentary time, with correlation coefficients ranging between -0.02 to 0.61 for single-item measures and -.02 to 0.49 for composite measures (16). Patterns of sedentary behavior, such as number and length of

breaks from sitting, are likely difficult to recall and better characterized using objective measures. However, domain-specific sedentary behaviors (e.g. time spent watching television, using the computer, and reading) are better captured by self-reports. Sorting out whether adverse health effects are due to sedentary behavior per se and/or to co-behaviors is best accomplished by studies which use both objective and self-reported measures.

While cohort studies should continue to collect self-report data, there is a need to add objective measures of sedentary behavior to these studies so as to increase understanding of whether and how sedentary behaviors impact health. This approach leverages ongoing cohort studies that have well-characterized participants and is cost saving compared to creating new, large cohort studies. Opportunities can leverage on-going prospective studies by adding objective activity measurement such as accelerometers and/or inclinometers depending on the definition of sedentary behavior researchers use (see proceedings from the first workshop for more discussion of definitions) (5). Previous studies have related self-reported sedentary behavior to biomarkers and clinical outcomes such as cardiovascular disease and diabetes (39), but few studies have used accelerometers or inclinometers, which can characterize sedentary behavior in more detail (e.g. sitting, standing, sit-to-stand transitions; bouts and durations of sedentary time). In adding objective measures to existing cohort studies, it appears likely that data analysis can be improved by use of newly emerging methods, such as iso-temporal substitutions (6). The method of iso-temporal substitution has advantages in determining whether sedentary behavior is associated with health outcomes independent of overall moderate-to-vigorous physical activity. Finally, data standardization methods would facilitate pooling results among studies (a recommendation from one of the other workshops) (5).

Repeated measures can provide information on patterns of sedentary behavior over time, and facilitate understanding the associations between sedentary behavior and health outcomes. For example, a recent study examined changes in self-reported sitting time over 2 years in community residing older adults (24). After adjustment for levels of moderate-to-vigorous physical activity, the study reported that those with excessive sedentary time at both baseline and 2 years had the highest rates of all-cause mortality\_(24). Replicating findings using studies with repeated objective measurements provides more compelling evidence that sustained sedentary behavior over time has adverse health outcomes. Adding repeated objective measures of sedentary time to existing studies could rectify evidence gaps related to hard clinical endpoints where almost no data exist (currently, data from studies of objectively measured sedentary behavior have focused primarily on biomarkers of disease risk).

Several cohort studies using objective measures are already underway. For example, participants in the Women's Health Study (anticipated n = 18,000) are currently wearing Actigraph (Pensacola, FL) accelerometers. Recognizing the potential limitations of accelerometers to measure posture (accelerometers cannot differentiate between sitting and standing with little motion), the Maastricht Study is using the activPAL™ inclinometer (PAL technologies Limited, Glasgow, United Kingdom) among approximately 10,000 adults with diabetes between ages 40 and 75 years. In contrast to accelerometers, inclinometers (such as activPAL™), can validly identify sitting/reclining separately from standing postures (22, 25). Collectively, these studies will offer opportunities to better understand measurement challenges and relate objectively derived sedentary behavior with health risk biomarkers and clinical health outcomes. Standardized methods for assessing sedentary behavior (and differentiation between sitting and standing) should be promoted so that results can be compared across studies (see Workshop 1 for

more discussion of standardization) (5). An ideal measure would involve 24 hour monitoring so that relationships among sedentary behavior, light activity, moderate to vigorous physical activity, napping, and night time sleep can be better understood; however, methods to identify each of these domains from the data output by monitors currently are not always well developed or standardized.

**Recommendation 2:** *Increase research efficiency by considering the advantages of a network of connected research studies and health systems such as: access to outcome and cost data from electronic health records (EHRs), population-based recruitment using EHRs to screen eligibility, and good ability to conduct pragmatic trials.*

**Rationale:** Many health systems have diverse and highly representative patient populations with clinical data captured in EHRs and robust research networks with experience in accessing, analyzing, and interpreting the data. For example, the Health Maintenance Organization (HMO) Research Network, a population-based research network consisting of 18 health care delivery organizations, constructed a virtual data warehouse of parallel databases housed at each site, with standard variable names and definitions. These databases can be used to construct analytic datasets across sites to conduct prospective and retrospective epidemiological studies. Health behaviors recorded in the EHR can be linked to morbidity and mortality data, health care costs, and health care utilization (20).

Health systems can provide opportunities to conduct targeted recruitment of high-risk individuals through information obtained through the EHR. Members can be recruited into trials or epidemiologic investigations for which outcomes can be passively ascertained through EHR data. Many health systems provide health care insurance to worksites, which can be targeted for

interventions, with outcomes from the interventions obtained through the EHR. Comparative effectiveness research and pragmatic, or real-world, trials can be conducted in health systems. These opportunities can provide research efficiencies in participant recruitment, outcome assessments, and identifying sites for interventions.

One current limitation is that health systems rarely systematically collect information on health behaviors such as physical activity and sedentary behavior in a manner to allow for data extraction from the EHR. This is beginning to shift and is anticipated to change over the next several years. For example, Kaiser Permanente has adopted an exercise vital sign (8) that routinely assesses patients' moderate to vigorous physical activity, and many Group Health Cooperative members complete annual health risk assessments that include items on self-reported sitting. While health behaviors are currently self-reported, wireless technologies are anticipated to be more fully utilized in health care in the future and could support the objective capture of health behaviors (4, 11). A current limitation may be the reluctance of health care providers to collect behavioral information, especially if it is not clear how the information would be used to improve patient care. Future studies could examine health care providers' views on collecting sedentary behavior information from their patients and providing advice to reduce sedentary behavior. An established risk of sedentary behavior on health outcomes may be needed prior to a systematized collection of patient sedentary behavior.

**Recommendation 3:** *Identify a variety of high-risk study populations and pre-plan collaboration among intervention studies so as to efficiently address the breadth of issues in sedentary behavior research including: effects of age and gender, dose-response, and effects of novel approaches (e.g. replace sedentary time with strength training or activities of different intensity).*

**Rationale:** In order to move the field of sedentary behavior research forward, studies targeting high-risk populations should be prioritized, as high-risk individuals can most benefit from sedentary behavior reduction. People who engage in more sedentary behavior may be more able to substantively decrease their sedentary behavior compared to increasing their physical activity (e.g. people with mobility limiting conditions, people healing from surgery). A threshold of sedentary behavior which connotes higher risk is not clear and, yet, there is a need to more thoroughly identify populations that spend excessive time doing sedentary behaviors. Objectively-assessed sedentary time from the US National Health and Nutrition Examination Survey show the top quartile of sedentary time to be 10.2 hours on average per day in a large sample of adults aged 20 to 59 years (31). Currently available guidelines recommend limiting screen time to less than two hours per day and to limit sedentary behavior in general (3, 19).

Current data suggest that adults over age 60 spend more time being sedentary than any other age group, on average approximately 8.5 hours per day (10, 29). Older adults with specific chronic conditions including breast cancer (26), prostate cancer (27), and heart failure (2), have been identified as having greater sedentary behavior than the general older adult population, although in some cases sample sizes are small. Older men are more sedentary than women and Hispanic older adults are less sedentary than White, Black, or other racial-ethnic groups (10). Sedentary

behavior has been related to physical function, falls, and other indicators of older adult frailty (7, 35).

Prevalence data using objective measures of sedentary behavior are currently incomplete, making it difficult to clearly elucidate populations that are most at-risk for engaging in high amounts of sedentary behavior. For example, it is not clear whether older adults in higher BMI categories are at higher risk for high sedentary behavior or whether there are multivariate risk profiles (e.g. older adults with low socioeconomic status and also diabetes and obesity). More detailed prevalence data using consistent measurement of sedentary behaviors are needed. In addition, most prevalence data include only total time spent engaging in sedentary behavior (measured by self-report or accelerometers) whereas the physiologic mechanisms of sedentary behavior suggest other outcomes may be as or more important—for example, the pattern of sedentary behavior accumulation throughout the day, sit-to-stand transitions, and bouts of sedentary behavior (17). Prevalence data could also include stratification by level of physical activity to help better understand whether there are protective effects. Potential high-risk populations could include frail older adults, postpartum women, people with mental illness, people with mobility disabilities or functional limitations, smokers, older adults living in assisted living, people with multiple chronic conditions such as cardiometabolic conditions and arthritis, and people undergoing life stage transitions (e.g. from childhood to adolescence, adolescence to young adulthood; retirement).

Interventions have started to target older adults with recognition that this group may be at elevated risk due to high sedentary behavior. There are two publications reporting on the short term (i.e. less than one month) feasibility of sedentary behavior reduction interventions in older adults which found reductions in sedentary behavior of ~2-3% (12, 14). The size of these

reductions may be statistically significant but it is not yet clear whether such changes are clinically meaningful. Studies have not yet elucidated the level of sedentary behavior reduction that connotes health benefits. Evidence for feasibility is, however, growing. A variety of different types of activities as substitution for sedentary behavior (e.g. strength training, high intensity training, standing) and different approaches to reducing sedentary behavior (e.g. reducing total time spent sedentary, reducing prolonged bouts of being sedentary, reducing certain types of sedentary behavior) need examination. Studies are also underway testing sedentary behavior reduction in people with type 2 diabetes (38) with one prior lifestyle based intervention in people with type 2 diabetes showing significant reductions in accelerometer measured sedentary behavior by 12 minutes per day at one year (9). Technology could be leveraged to support these sedentary behavior interventions, although some high-risk populations may have barriers to using technology (33, 40).

**Recommendation 4:** *In intervention research, include comparative-effectiveness designs (specifically comparisons of interventions to reduce sedentary behavior versus interventions to increase moderate to-vigorous physical activity) and pure environmental interventions, so as to enhance later efforts to translate interventions into practice. Studies should also address factors other than efficacy that can affect translation such as reach, cost, adverse effects, adherence, and sustainability.*

**Rationale:** Comparative effectiveness research (CER) “informs health-care decisions by providing evidence on the effectiveness, benefits, and harms of different treatment options”(1). CER trials could compare sedentary behavior reduction to increased moderate-to-vigorous physical activity. Some effects might be similar (e.g. depressive symptoms) but others might be

unique (e.g. cardiovascular fitness might be more favorably changed in the moderate to vigorous physical activity group while general mobility might be more favorably changed through sedentary behavior reduction). Some people might benefit or adhere more to physical activity while for others it may be sedentary behavior reduction. CER and associated study designs (e.g. Sequential Multiple Assignment Randomized Trials) (23) can help the field better understand these differences and identify the best intervention for a particular person given their preferences, health status, and life situation.

In addition, comparative effectiveness trials could compare different approaches to sedentary behavior reduction. Frieden's health impact pyramid suggests that the largest impact occurs by "changing the context to make individuals' default decisions healthy"(13). Purely environmental interventions could be compared to multilevel interventions (following the logic of the ecological model which includes individual, interpersonal, and built environment changes) (32). To-date, the few published interventions have incorporated some multilevel approaches such as enhancing motivation and addressing social and environmental constraints. For example, one workplace intervention consisted of individual (health coaching consultation sessions, self-regulation strategies, and motivational interviewing), environmental (sit-stand workstations), and organizational support approaches to reduce sitting time by 125 minutes per day over 4 weeks (18). A lifestyle, community-based intervention in people with diabetes used various behavioral modification strategies to reduce self-reported sitting by 30 minutes per day over one year and 12 minutes by accelerometers (9). Some purely environmental interventions have been conducted as well. Pronk et al. provided sit-stand workstations to employees with sedentary jobs and found a 66 minute reduction in sitting over 4 weeks; the effect disappeared after the workstations were removed (34). One home-based study provided television lock out devices to overweight and

obese adults and, in comparison to a control group, found trends for increased energy expenditure over 3 weeks (30). These studies suggest a slightly larger effect for multilevel approaches, but studies were short in duration, had incomparable measurement methods, and study designs differed. Larger trials that can compare different approaches (e.g. purely environmental versus multilevel; physical activity versus sedentary behavior) and assess important unstudied issues such as reach, cost, adherence, and sustainability.

There is also a need to examine different strategies for changing sedentary behavior and the impact of different types of intervention goals. For example, researchers could compare the effects of different messaging strategies (e.g. comparing messages to “sit less” versus “move more;” also discussed in another workshop) (28). In addition, better understanding which of these types of goals has the best adherence (or whether certain types of people respond best to different types of goals) will help in translating controlled research findings into real world interventions. Finally, qualitative information is vital in understanding causes of excessive sedentary behavior. This information is needed to help researchers understand barriers, beliefs, and attitudes around reducing sedentary behavior and acceptability of different intervention and measurement approaches.

## **Conclusions**

There are opportunities to gain research efficiencies by leveraging existing epidemiologic cohorts and health systems. Existing and new studies can move towards capturing sedentary behavior with objective monitoring, instead of relying on self-reports, so that patterns of sedentary behavior and relationships with longer term hard health outcomes can be elucidated. Health systems can provide an excellent setting for pragmatic trials and observational studies examining relationships of sedentary behavior with health outcomes, health costs, and utilization. Finally, a variety of interventions targeting high-risk groups and CER approaches need to be undertaken to move the field of research forward in an inclusive manner.

## **Acknowledgements**

The workshop was sponsored by the National Heart, Lung and Blood Institute, the National Institute on Aging and the Office of Disease Prevention of the National Institute of Health.

The views expressed in this document reflect the collective ideas and opinions of the authors and does not necessarily represent the official views of the National Institutes of Health , or the U.S. Department of Health and Human Services and the other workshop participants. Support for this work provided in part by a Shahid & Ann Carlson Khan Professorship (Dr. Buchner), grant CA154647 from the US National Institutes of Health (Dr. Lee, by National Health and Medical Research Council of Australia Program Grant (#569940) and Research Fellowship funding (#1003960) and by Victorian Government OIS funding (Dr. Owen). Authors were provided with an honorarium from the National Institutes of Health for their participation in the workshop. We wish to thank the discussants for this workshop: Dr. Stephen P. Fortmann, Kaiser Permanente Center for Health Research, Dr. James F. Sallis, University of California San Diego, Dr. William Haskell, Stanford University School of Medicine, and Dr. Steven N. Blair, University of South Carolina. The results of the present study do not constitute endorsement by ACSM.

**Conflict of Interest:** None

## References

1. Agency for Healthcare Research and Quality [Internet]. Washington D.C. Available from: <http://effectivehealthcare.ahrq.gov/index.cfm/what-is-comparative-effectiveness-research1/>.
2. Alosco ML, Spitznagel MB, Miller L et al. Depression is associated with reduced physical activity in persons with heart failure. *Health Psychol.* 2012;31(6):754-62.
3. American Cancer Society [Internet]. American Cancer Society; [cited 2014 June 19]. Available from: <http://www.cancer.org/acs/groups/cid/documents/webcontent/002577-pdf.pdf>.
4. Avancha S, Baxi A. Privacy in mobile technology for personal healthcare. *ACM Computing Surveys.* 2012;45(1):3:1-3:54.
5. Barone Gibbs B, Hergenroeder A, Katsmarzyk P, Lee I, Jakicic JM. Definition, measurement and health risks associated with sedentary behavior in adults. In. *Med Sci Sports Exerc.* In press.
6. Buman MP, Winkler EA, Kurka JM et al. Reallocating time to sleep, sedentary behaviors, or active behaviors: associations with cardiovascular disease risk biomarkers, NHANES 2005-2006. *Am J Epidemiol.* 2014;179(3):323-34.
7. Cawthon PM, Blackwell TL, Cauley JA et al. Objective assessment of activity, energy expenditure, and functional limitations in older men: the Osteoporotic Fractures in Men study. *J Gerontol A Biol Sci Med Sci.* 2013;68(12):1518-24.

8. Coleman KJ, Ngor E, Reynolds K et al. Initial validation of an exercise "vital sign" in electronic medical records. *Med Sci Sports Exerc.* 2012;44(11):2071-6.
9. De Greef KP, Deforche BI, Ruige JB et al. The effects of a pedometer-based behavioral modification program with telephone support on physical activity and sedentary behavior in type 2 diabetes patients. *Patient Educ Couns.* 2011;84(2):275-9.
10. Evenson KR, Buchner DM, Morland KB. Objective measurement of physical activity and sedentary behavior among US adults aged 60 years or older. *Prev Chronic Dis.* 2012;9:E26.
11. Fiordelli M, Diviani N, Schulz PJ. Mapping mHealth research: a decade of evolution. *J Med Internet Res.* 2013;15(5):e95.
12. Fitzsimons CF, Kirk A, Baker G, Michie F, Kane C, Mutrie N. Using an individualised consultation and activPAL feedback to reduce sedentary time in older Scottish adults: results of a feasibility and pilot study. *Prev Med.* 2013 57(5):718-20.
13. Frieden TR. A framework for public health action: the health impact pyramid. *Am J Public Health.* 2010;100(4):590-5.
14. Gardiner PA, Eakin EG, Healy GN, Owen N. Feasibility of reducing older adults' sedentary time. *Am. J. Prev. Med.* 2011;41(2):174-7.
15. Gardiner PA, Healy GN, Eakin EG et al. Associations between television viewing time and overall sitting time with the metabolic syndrome in older men and women: the Australian Diabetes, Obesity and Lifestyle study. *J Am Geriatr Soc.* 2011;59(5):788-96.

16. Healy GN, Clark BK, Winkler EA, Gardiner PA, Brown WJ, Matthews CE. Measurement of adults' sedentary time in population-based studies. *Am J Prev Med.* 2011;41(2):216-27.
17. Healy GN, Dunstan DW, Salmon J et al. Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care.* 2008;31(4):661-6.
18. Healy GN, Eakin EG, Lamontagne AD et al. Reducing sitting time in office workers: short-term efficacy of a multicomponent intervention. *Prev Med.* 2013;57(1):43-8.
19. Heart Foundation [Internet]. National Heart Foundation of Australia; [cited 2014 June 19]. Available from: <http://www.heartfoundation.org.au/SiteCollectionDocuments/HW-PA-SittingLess-Adults.pdf>.
20. HMO Research Network [Internet]. Seattle, WA: HMO Research Network. Available from: [http://www.hmoresearchnetwork.org/resources/toolkit/HMORN\\_CollaborationToolkit.pdf](http://www.hmoresearchnetwork.org/resources/toolkit/HMORN_CollaborationToolkit.pdf).
21. Howard BJ, Balkau B, Thorp AA et al. Associations of overall sitting time and TV viewing time with fibrinogen and C reactive protein: the AusDiab study. *Br J Sports Med.* 2014 Feb 18 [ePub Ahead of Print].
22. Kozey-Keadle S, Libertine A, Lyden K, Staudenmayer J, Freedson PS. Validation of wearable monitors for assessing sedentary behavior. *Med Sci Sports Exerc.* 2011;43(8):1561-7.
23. Lei H, Nahum-Shani I, Lynch K, Oslin D, Murphy SA. A "SMART" design for building individualized treatment sequences. *Annu Rev Clin Psychol.* 2012;8:21-48.

24. Leon-Munoz LM, Martinez-Gomez D, Balboa-Castillo T, Lopez-Garcia E, Guallar-Castillon P, Rodriguez-Artalejo F. Continued sedentariness, change in sitting time, and mortality in older adults. *Med Sci Sports Exerc.* 2013;45(8):1501-7.
25. Lyden K, Kozey-Keadle SL, Staudenmayer JW, Freedson PS. Validity of two wearable monitors to estimate breaks from sedentary time. *Med Sci Sports Exerc.* 2012 44(11):2243-52.
26. Lynch BM, Dunstan DW, Healy GN, Winkler E, Eakin E, Owen N. Objectively measured physical activity and sedentary time of breast cancer survivors, and associations with adiposity: findings from NHANES (2003-2006). *Cancer Causes Control.* 2010;21(2):283-8.
27. Lynch BM, Dunstan DW, Winkler E, Healy GN, Eakin E, Owen N. Objectively assessed physical activity, sedentary time and waist circumference among prostate cancer survivors: findings from the National Health and Nutrition Examination Survey (2003-2006). *Eur J Cancer Care (Engl).* 2011;20(4):514-9.
28. Manini TM, King AC, Marshall SJ, Robinson TN, Rejeski WJ. Interventions to reduce sedentary behavior. In. *Med Sci Sports Exerc.* In press.
29. Matthews CE, Chen KY, Freedson PS et al. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol.* 2008;167(7):875-81.
30. Otten JJ, Jones KE, Littenberg B, Harvey-Berino J. Effects of television viewing reduction on energy intake and expenditure in overweight and obese adults: a randomized controlled trial. *Arch Intern Med.* 2009;169(22):2109-15.

31. Owen N, Sparling PB, Healy GN, Dunstan DW, Matthews CE. Sedentary behavior: emerging evidence for a new health risk. *Mayo Clin Proc.* 2010;85(12):1138-41.
32. Owen N, Sugiyama T, Eakin EE, Gardiner PA, Tremblay MS, Sallis JF. Adults' sedentary behavior determinants and interventions. *Am J Prev Med.* 2011;41(2):189-96.
33. Parker SJ, Jessel S, Richardson JE, Reid MC. Older adults are mobile too! Identifying the barriers and facilitators to older adults' use of mHealth for pain management. *BMC Geriatr.* 2013;13:43.
34. Pronk NP, Katz AS, Lowry M, Payfer JR. Reducing occupational sitting time and improving worker health: the Take-a-Stand Project, 2011. *Prev Chronic Dis.* 2012;9:E154.
35. Seguin R, Lamonte M, Tinker L et al. Sedentary Behavior and Physical Function Decline in Older Women: Findings from the Women's Health Initiative. *J Aging Res.* 2012;2012:271589.
36. Sidney S, Sternfeld B, Haskell WL, Jacobs DR, Jr., Chesney MA, Hulley SB. Television viewing and cardiovascular risk factors in young adults: the CARDIA study. *Ann Epidemiol.* 1996;6(2):154-9.
37. Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med.* 2011;41(2):207-15.

38. Wilmot EG, Davies MJ, Edwardson CL et al. Rationale and study design for a randomised controlled trial to reduce sedentary time in adults at risk of Type 2 Diabetes Mellitus: Project STAND (Sedentary Time AND Diabetes). *BMC Public Health*. 2011;11(1):908.
39. Wilmot EG, Edwardson CL, Achana FA et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia*. 2012;55(11):2895-905.
40. Young R, Willis E, Cameron G, Geana M. "Willing but unwilling": attitudinal barriers to adoption of home-based health information technology among older adults. *Health informatics journal*. 2014;20(2): 127-135.