

Medicine & Science IN Sports & Exercise

The Official Journal of the American College of Sports Medicine
www.acsm-msse.org

. . . Published ahead of Print

Sedentary Behavior Research Priorities—NHLBI/NIA Sedentary Behavior Workshop Summary

Josephine Boyington¹, Lyndon Joseph², Roger Fielding³, and Russell Pate⁴

¹National Heart, Lung and Blood Institute, National Institutes of Health, Bethesda, MD;

²National Institute on Aging, National Institutes of Health, Bethesda, MD; ³Nutrition, Exercise Physiology, and Sarcopenia Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA; ⁴Arnold School of Public Health, University of South Carolina, Columbia, SC

Accepted for Publication: 28 July 2014

Medicine & Science in Sports & Exercise® **Published ahead of Print** contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Sedentary Behavior Research Priorities—NHLBI/NIA Sedentary Behavior Workshop Summary

Josephine Boyington¹, Lyndon Joseph², Roger Fielding³, and Russell Pate⁴

¹National Heart, Lung and Blood Institute, National Institutes of Health, Bethesda, MD;

²National Institute on Aging, National Institutes of Health, Bethesda, MD; ³Nutrition, Exercise Physiology, and Sarcopenia Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA; ⁴Arnold School of Public Health, University of South Carolina, Columbia, SC

Corresponding Author:

Russell Pate, Ph.D.

Arnold School of Public Health, University of South Carolina

Suite 212, Public Health Research Center

921 Assembly Street

Columbia, SC 29208

Phone: (803) 777-2456

Email: rpate@mailbox.sc.edu

Running Title: Sedentary Behavior Research Priorities

Acknowledgments:

The workshop was sponsored by the National Heart, Lung and Blood Institute, the National Institute on Aging and the Office of Disease Prevention of the National Institutes of Health. The views expressed in this document reflect the collective ideas and opinions of the authors and do not necessarily represent the official views of the National Institutes of Health, the U.S. Department of Health and Human Services or the other workshop participants.

Dr. Fielding's participation in this project was partially supported by the U.S. Department of Agriculture, under agreement No. 58-19500-014, and the Boston Claude D. Pepper Older Americans Independence Center (1P30AG031679). Any opinions, findings, conclusion, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Agriculture.

Conflicts:

Dr. Fielding receives grant support from the USDA, National Institutes of Health, Nestec Ltd. and Regeneron Pharmaceutical; is a consultant to Regeneron Pharmaceuticals and Nestec, Ltd; serves on the Scientific Advisory Boards of Cytokinetics Inc, Pronutria Inc. Amonnett, Inc., and Inside Tracker Inc,; and owns stock or options in Pronutria, Amonnett Inc., Inside Tracker, and Myosyntax.

Dr. Pate receives grant support from the National Institutes of Health, Centers for Disease Control and Prevention, Duke Endowment, and FTZ Coca-Cola Service Company. He serves on the Scientific Advisory Board of Curves, Inc. The other authors have no potential conflicts.

Introduction and Workshop Purpose

Over the course of 2013, the National Heart, Lung and Blood Institute (NHLBI) and the National Institute on Aging (NIA) collaborated with a multi-national group of expert scientists to convene, host and facilitate a four-part virtual webinar workshop on sedentary behavior. Participants met to evaluate and discuss the current epidemiological, clinical and other relevant literature on sedentary behavior and to identify research priorities relative to preventing cardiovascular disease and mitigating aging-related disability and functional outcomes. The workshop was predicated on findings of insufficient levels of population adherence to current physical activity recommendations (1,12) and the need to address the NHLBI and NIA research portfolio gaps in the area of sedentary behavior.

Scientific Rationale

No consistent definition of sedentary behavior currently exists. However, sitting, lying down, watching television, and participating in other screen-based activities are collectively labeled sedentary behaviors. In general, sedentary behavior consists of activities that do not increase energy expenditure markedly above resting metabolic levels (15). Consequently, they qualify as low energy expenditure activities, (15) which are associated with adverse metabolic and physical health outcomes with prolonged exposure (5,7-11,18). In contrast, high energy expenditure activities confer health benefits with prolonged exposure, and this knowledge undergirds current physical activity recommendations (2,6,17,19). Although physical activity recommendations for moderate-to-vigorous intensity activities are based on strong evidence, they have yet to translate into sustained changes in population-level behaviors. In fact, the majority of the US population is considered mostly inactive and sedentary (2,3,6).

Yet, the health consequences of sedentary behavior are not clearly understood (4,5,7-11,18). Current knowledge suggests negative associations between sedentary behavior and health outcomes and indicates that the adverse effects associated with sedentary behavior are separate from those associated with low levels of moderate-to-vigorous intensity activities (13,14). Consequently, a knowledge gap remains regarding the independent effects of sedentary behavior on health outcomes, how sedentary behavior research fits into the vast knowledge base on physical activity, and how current and future knowledge about sedentary behavior can inform future health recommendations.

Framing the Workshop

In an effort to engage all stakeholders, the NHLBI and NIA released a joint “Request for Information (RFI)” in January 2013 entitled “Request for Information (RFI) for High-Priority Research Topics for Sedentary Behavior - NOT-HL-13-166” (16). This RFI sought stakeholders’ perspectives on unanswered research questions in several areas, including approaches needed to evaluate and measure sedentary behavior in different populations, types of interventions and strategies needed to reduce the high prevalence of sedentary behaviors in the U.S., and identification of barriers and opportunities for translating current scientific knowledge into population health improvements. The substantial number of responses NHLBI and NIA received guided the workshop planning, identification of discussion topics and formulation of the four thematic sessions: 1. Epidemiology of Sedentary Behavior; 2. Physiology of Sedentary Behavior and its Relationship to Health Outcomes; 3. Influences on Sedentary Behavior and Interventions to Reduce Sedentary Behavior; and 4. Novel Strategies for Sedentary Behavior Research.

Pre-meeting Activities and Charge to the Key Participants

Leading expert scientists in the four thematic areas were invited to participate as session coordinators, key presenters or audience participants. Teams of five to seven experts developed four or five concise 10-minute presentations, which served as the framework for discussions in each of the four, 2.5-hour virtual meetings. An additional group of invited scientists served as panel discussants for each session and provided summary comments and provocative questions that fostered deeper discussion of key issues. All of the discussions were facilitated by the meeting co-chairs, Dr. Russell Pate and Dr. Roger Fielding.

Workshop Summary

To capture the key findings of the workshop, the meeting chair concluded each session by identifying the two top priority recommendations. Comprehensive summaries of the meetings are presented in four brief white papers that accompany this document, and include the substance of the presentations and discussions, a list of all the recommendations that emerged, and the final two priority recommendations for each session. Each of the white papers also reflects the specific themes discussed in one of the four sessions, and includes substantial input from the session expert scientists and discussants convened for that particular session. As such, the authors of the white papers had the flexibility to present their recommendations in the manner that best captured their sessions' theme, tone and discussion. Across all sessions, participants varied in their use and definition of the concept of sedentary behavior. In fact the term 'sedentary behavior', 'sedentary time', and 'sedentariness' were all used to inform the discussions. Consequently, in the interest of authenticity, the four white papers also reflect this variety of usage and perspectives. In the paragraphs below,

summaries of the discussions for each of the workshop sessions are presented briefly. However, details with supporting rationale for all the recommendations can be found in the accompanying white papers.

Webinar #1 – The Epidemiology of Sedentary Behavior

The initial workshop session was dedicated to reviewing and discussing several issues basic to the study of sedentary behavior and its impact on health. Much attention was given to definitions of sedentary behavior proposed previously and their implications for measurement of sedentary behavior in research studies. Workshop participants reviewed the current status of epidemiological research that has examined relationships between sedentary behavior and health outcomes and overviewed studies that have examined interactive influences of sedentary behavior and physical activity. They concluded that consensus regarding conceptual and operational definitions of sedentary behavior, better methods for measuring sedentary behavior in research studies, and large-scale prospective observational studies of the effects of sedentary behavior on multiple health outcomes are needed.

Webinar # 2 – Physiology of Sedentary Behavior and its Relationship to Health Outcomes

The second webinar focused on the current state of understanding of the biological/physiological mechanisms that underlie sedentary behavior. Key areas of focus included cardio-metabolic consequences of sedentary behavior, animal models of sedentary behavior or forced inactivity, and the interaction between central neurologic drivers of sedentary behavior and peripheral adaptation and feedback. Limitations in current understanding of the molecular changes that occur with sedentary behavior were noted, in particular the fact that the molecular evidence for

sedentary behavior effects is derived primarily from studies of conditions and models of low physical activity. Participants resolved that while low physical activity and sedentary behavior are likely to be related mechanistically, they are distinct entities. Discussion also focused on the need to develop better methodologies to address sedentary behavior, including further investigation of appropriate animal models and systems. Webinar participants identified several key research needs going forward, including understanding the interaction between central and peripheral mediators of sedentary behavior and the resultant biological consequences, understanding the molecular basis for the role of changes in sedentary behavior in accelerating the loss of aerobic capacity, and examining muscle performance with advancing age.

Webinar # 3 – Influences on Sedentary Behavior/Interventions to Reduce Sedentary

Behavior

The third webinar focused on intervention strategies designed to reduce sedentary behavior. The main conclusion from this webinar was that interventions targeting sedentary behavior are conceptually and operationally distinct from interventions targeting physical activity. In reviewing the evidence on at-risk populations (e.g., children, older adults) and levels of intervention (e.g., community, workplace, individual), participants concluded that the needs of populations vary and recommended that future endeavors carefully consider the levels (community, work place, individual) for which interventions are to be developed and the populations to which they are targeted . The complexity of messaging to the public and the biomedical community surrounding the topic of sedentary behavior and a clear need for consensus around this message also were considered. Finally, participants proposed novel approaches to address questions related to the dose-response effects of changing sedentary

behavior, the establishment of clinically-meaningful changes in sedentary behavior, and development of more sophisticated tools to quantify sedentary behavior exposure.

Webinar #4 – Novel Strategies for Sedentary Behavior Research

The final workshop session focused on unique and efficient strategies for conducting research on sedentary behavior and health. The workshop examined the need for research on sedentary behavior across a wide range of issues and noted that most of the current evidence linking sedentary behavior to health is based on observational studies. Ultimately, evidence will be needed to establish public health guidelines and public health interventions aimed at reducing sedentary behavior. Much discussion focused on the availability of databases that could support important research on sedentary behavior. These include datasets generated in ongoing large-scale epidemiologic studies in which accelerometry data have been or are being collected. These data, although typically collected for the purpose of measuring physical activity, can be reduced to measure sedentary behavior as well, and thus are a potentially valuable resource for future research in this area. Likewise, patient data included in the electronic health records of large integrated health systems provide opportunities for comparative effectiveness research and observational studies of sedentary behavior in patient populations.

Conclusion

Paragraph 10: This workshop was convened to engage expert extramural scientists in a series of discussions to identify critical gaps in the literature on sedentary behavior and cardiovascular and aging health outcomes. Themes that cut across the four meeting sessions included the need to develop a consensus definition, validate and standardize assessment methods (both objective and subjective) to provide for accuracy and consistency, and develop appropriate scientific models of research. Other topics included the use of translational models to understand personal, social and environmental determinants of sedentary behavior; development of pragmatic trials using large networks of integrated health systems to aid with measurement, prevalence and incidence data collection; exploring the link between incidence rate and risk factors; and the use of technology. Significant discussions focused on whether sedentary behavior should be considered as a distinct and separate behavior from physical inactivity with separate adverse effects on health. To this end, the workshop summary proposed that future studies should seek to examine the separate and interactive effects of sedentary behavior and moderate-to-vigorous physical activity. In addition, it highlighted the importance of looking at patterns of sedentary behavior, developing and testing multi-level interventions, and exploring dose-response relationships of sedentary behavior to health outcomes. Participants acknowledged that sedentary behavior presents a unique challenge for which the strategies employed to increase physical activity may not be suited. That notwithstanding, the research community should make concerted efforts to study and understand sedentary behaviors and to intervene to effect positive health outcomes and ensure that appropriate resources be directed to this endeavor.

Acknowledgments:

The workshop was sponsored by the National Heart, Lung and Blood Institute, the National Institute on Aging and the Office of Disease Prevention of the National Institutes of Health. The views expressed in this document reflect the collective ideas and opinions of the authors and do not necessarily represent the official views of the National Institutes of Health, the U.S. Department of Health and Human Services or the other workshop participants.

Dr. Fielding's participation in this project was partially supported by the U.S. Department of Agriculture, under agreement No. 58-19500-014, and the Boston Claude D. Pepper Older Americans Independence Center (1P30AG031679). Any opinions, findings, conclusion, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Agriculture.

Conflicts:

Dr. Fielding receives grant support from the USDA, National Institutes of Health, Nestec Ltd. and Regeneron Pharmaceutical; is a consultant to Regeneron Pharmaceuticals and Nestec, Ltd; serves on the Scientific Advisory Boards of Cytokinetics Inc, Pronutria Inc. Amonnett, Inc., and Inside Tracker Inc.,; and owns stock or options in Pronutria, Amonnett Inc., Inside Tracker, and Myosyntax.

Dr. Pate receives grant support from the National Institutes of Health, Centers for Disease Control and Prevention, Duke Endowment, and FTZ Coca-Cola Service Company. He serves on the Scientific Advisory Board of Curves, Inc.

The other authors have no potential conflicts.

References:

1. Blair SN, Kohl HW, III, Paffenbarger RS, Jr., Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA*. 1989;262(17):2395-401.
2. Centers for Disease Control and Prevention Web site [internet]. Atlanta (GA): One in five adults meet overall physical activity guidelines; [cited 2013 September 1]. Available from: <http://www.cdc.gov/media/releases/2013/p0502-physical-activity.html>.
3. Cohen SS, Matthews CE, Signorello LB, Schlundt DG, Blot WJ, Buchowski MS. Sedentary and physically active behavior patterns among low-income African-American and white adults living in the southeastern United States. *PLoS One*. [Internet] 2013 [cited 2013 September 1]; 8(4):e59975. Available at <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0059975>. doi:10.1371/journal.pone.0059975
4. Evenson KR, Buchner DM, Morland KB. Objective measurement of physical activity and sedentary behavior among US adults aged 60 years or older. *Prev Chronic Dis*. [Internet] 2012 [cited 2013 September 1]; 9:110109. Available from http://www.cdc.gov/pcd/issues/2012/11_0109.htm. doi:<http://dx.doi.org/10.5888/pcd9.110109>
5. Ford ES, Kohl HW III, Mokdad AH, Ajani UA. Sedentary behavior, physical activity, and the metabolic syndrome among U.S. adults. *Obes Res*. 2005;13(3):608–614.
6. Haskell WL. J.B. Wolffe Memorial Lecture. Health consequences of physical activity: understanding and challenges regarding dose-response. *Med.Sci.Sports Exerc*. 1994;26(6):649-60.

7. Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, Rimm EB. Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med.* 2001;161(12): 1542–1548.
8. Hu FB, Li TY, Colditz GA, Willett WC, Manson JE. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *JAMA.* 2003; 289(14):1785–1791.
9. Jakes RW, Day NE, Khaw KT, Luben R, Oakes S, Welch A, Bingham S, Wareham NJ. Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. *Eur J Clin Nutr.* 2003; 57(9): 1089–1096.
10. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc.* 2009; 41(5): 998–1005.
11. LeBlanc AG, Spence JC, Carson V, Connor Gorber S, Dillman C, Janssen I, Kho ME, Stearns JA, Timmons BW, Tremblay MS. Systematic review of sedentary behaviour and health indicators in the early years. (aged 0-4 years). *Appl Physiol Nutr Metab.* 2012;37(4):753-72.
12. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, Troiano RP. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol.* 2008; 167(7):875-81.
13. Matthews CE, George SM, Moore SC, Bowles HR, Blair A, Park Y, Troiano RP, Hollenbeck A, Schatzkin A. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr.* 2012;95(2):437-45.

14. Owen N, Sparling PB, Healy GN, Dunstan DW, Matthews CE. Sedentary behavior: emerging evidence for a new health risk. *Mayo Clin Proc.* 2010;85(12):1138-41.
15. Pate RR, O'Neill JR, Lobelo F. The evolving definition of "sedentary". *Exerc Sport Sci Rev.* 2008;36(4):173-8.
16. Request for Information (RFI) for High-Priority Research Topics for Sedentary Behavior Notice Number: NOT-HL-13-166. Available from: <http://grants.nih.gov/grants/guide/notice-files/NOT-HL-13-166.html>
17. Shaper AG, Wannamethee G, Weatherall R. Physical activity and ischaemic heart disease in middle-aged British men. *Br.Heart J.* 1991;66(5):384-94.
18. Tremblay MS, LeBlanc AG, Kho ME, Saunders TJ, Larouche R, Colley RC, Goldfield G, Connor Gorber S. Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act.* [Internet]. 2011 [cited 2013 September 1];8:98. Available from: <http://www.ijbnpa.org/content/8/1/98>. doi:10.1186/1479-5868-8-98.
19. U.S. Department of Health and Human Services. *Physical Activity Guidelines Advisory Committee Report, 2008*. U.S. Department of Health and Human Services, Rockville, MD: Office of the Secretary. Available from: U.S. GPO, Washington.